

14. (Amended) The method according to claim 8, wherein the detection reagent is an antibody or immunoreactive fragment thereof.

15. (Amended) The method according to claim 8, wherein the detection reagent is labelled by a fluorophore or chromophore.

16. (Amended) The method according to claim 7, wherein the specific binding pair is biotin-avidin or biotin-streptavidin.

17. (Amended) The method according to claim 1, wherein the sample is an undiluted serum sample.

18. (Amended) The method according to claim 1, wherein the sample is an undiluted whole blood sample.

20. (Amended) The test kit according to claim 19, wherein the ratio between the receptor-binding capacity of ligand immobilized on the solid phase and the ligand-binding capacity of the analyte-specific receptor substance is in the range of from about 1:2 to about 1:1000.

21. (Amended) The test kit according to claim 19 or 20, further comprising a lateral flow membrane strip having said receptor-binding ligand immobilized in or on a reaction zone of the membrane and having said analyte-binding receptor substance dissolvably pre-deposited in or on the membrane upstream of the reaction zone.

23. (Amended) The test kit according to claim 22, wherein the ratio between the amount of ligand-binding analyte-specific receptor and the total amount of analyte-specific receptor is in the range of from about 1:2 to about 1:1000.

24. (Amended) The test kit according to claim 22 or 23, further comprising a lateral flow membrane strip having said receptor-binding ligand immobilized in or on a reaction zone of the membrane and having said analyte-binding receptor substance dissolvably pre-deposited in or on the membrane upstream of the reaction zone.

26. (Amended) The test kit according to claim 25, wherein the ratio between said second amount of analyte-binding receptor substance immobilized to the solid phase, and the sum of said first

Please add the following new claims.

--29. The method according to claim 9, wherein the ratio between said isolated fraction of the amount of active analyte-binding receptor and the total amount of active analyte-binding receptor contacted with the sample is in the range of from about 1:5 to 1:100.

30. The method according to claim 9, wherein the ratio between said isolated fraction of the amount of active analyte-binding receptor and the total amount of active analyte-binding receptor contacted with the sample is no more than about 1:20.

31. The method according to claim 10, wherein said flow matrix is a lateral flow matrix.

32. The method according to claim 29, wherein said lateral flow matrix is a membrane strip.

33. The test kit according to claim 20, wherein the ratio between the receptor-binding capacity of ligand immobilized on the